GENERAL INDEX TO VOLUME XXXIV

New scientific names of plants and the final members of new combinations are printed in bold face type; synonyms and page numbers having reference to figures and plates, in italics; and previously published names and all other matter, in ordinary type.

Adiantum pedatum, 122 Adlumia, 191; fungosa, 188, 239 Alkaloidal properties of Corydalis, 195 American origin of the cultivated Cucurbits, 101

Anderson, Edgar: Field studies of Guatemalan maize, 433; William L. Brown and. The northern flint corns, 1

Andrews, Henry N.: John Henry Brittsphysician and fossil hunter, 115; and Ellen M. Kern. The Idaho Tempskyas and associated fossil plants, 119; and Lee W. Lenz. Fossil Polypores from Idaho, 113

Anemia elegans, 149; Fremonti, 152 Antigua, Guatemala, typical ear of corn from region about, 452, 456

Archeological maize in the United States, 8, 10, 13, 28

Asclepias incarnata, 356; speciosa × syriaca, 358; syriaca, 355, 358; tuberosa, 353, cl. interior, 353, ssp. interior, 353, 361, intraspecific differentiation of, 409, ssp. Rolfsii, 361, ssp. tuberosa, 361

Asclepias tuberosa, Some dynamics of leaf variation in, 353

Asclepias tuberosa: biology of, 354; cytogenetics of, 358, 359; ecology of, 359; hybridization in, 391; leaf types of three subspecies, 361, 366, 367; phenocontour maps, 370; population studies, 362; reproductive habit, 355; roadside profile of, 394, 395, 401; compared with herbarium profile, 403, 407; vegetative habit,

Austroclepis australis, 146

Bennettiales, fossil, 153 Bolivia, wild rubber in, 266 Borckhausenia, 197 Brettanomyces, 33 Britts, John Henry-physician and fossil hunter, 115, 115 Brown, William L., and Edgar Anderson.

The northern flint corns, 1 Bulbocapnos, 197, 207

Burkholder's medium, growth of yeast cultures on, 76, 81, 85, 95 Butterflyweed, see Asclepias tuberosa

Candida Guilliermondii, 33; mortifera, 33 Caoutchoua elastica, 292 Capnites, 197, 207

Capnodes, 197; aureum, 229; Bidwellianum, 201; Caseanum, 201; crystallinum, 217; flavulum, 215; pauciflorum, 207; Scouleri, 199

Capnoides, 188, 197, 209; brachycarpum, 204; Brandegei, 203; campestre, 222; curvisiliquum, 223, 226; Cusickii, 205; Engelmannii, 230; euchlamydeum, 230; glauca, 211; Halei, 217, 222, 223; bastatum, 206; macrorrbiza, 230; micranthum, 219; montanum, 234; pachylobum, 234; pauciflorum, 207; Scouleri, 199; sempervirens, 188, 211; Wetherillii, 230

Carnation (Dianthus caryophyllus), Inheritance in the, III. Inheritance of flower color, 39, 72, 74

Cells, relation between viability of, and inability to grow on deficient medium, 83

Ceratopteris thalictroides, 149

Chimaltenango, Guatemala, corn from, 462 Chromosomal complements of Corydalis, 194

Chromosome knob numbers in dent corn inbreds, 5

Cibotium sp., compared with Tempskya Wessellii, 133

Cisticapnos, 188

Clepsydropsis australis, 146

Clons of butterflyweed, 353, 354

Coal mines, fossil plants of, collected by Dr. Britts, 115

Color in carnations, inheritance of, 39, 72, 74; acyanic group, 41; cyanic group, 48; transition group, 56; variegated group,

Coniferales, fossil, 153

Corn: archeology of, 10; breeding, 16, 449; brewing varieties, 445; dent varieties, 1, 15; field studies of, in Guatemala, 433; hybridization of, 15; Mexican, 14; "mul-

19

Flo

Fo

Fo

Fo

Foe

Fu

Ga

Gei

Ger

Geo

Gle

Go

Gu

Her

Hei

Hei

Hev

S

S

p

C

P

b

3

2

3

3

a

Hev

tiplication" in Guatemalan varieties, 448; northern flint, 1; of the Corn Belt, 14; popping varieties, 445; prehistoric collections, 9, 28

Corydalis: Monograph of the North American species of, 187; morphology of, 188, of flowers, 252-258; taxonomy, 197

Corydalis, 187; Albertae, 229; Allenii, 199; annua, 211; aurea ssp. aurea, 229, ssp. occidentalis, 234; aurea var. australis, 222, \$ crystallina, 217, var. curvisiliqua, 226, a flavula, 215, var. macrantha, 229, var. micrantha, 219, var. occidentalis, 234, var. parviflora, 229, var. robusta, 229, var. typica, 229; biaurita, 239; Bidwelliae, 201; bilimbata, 234; brachycarpa, 204; bracteosa, 239; Brandegei, 203; bulbosa, 194; campestris, 222; canadensis, 239; Caseana, 201, ssp. brachycarpa, 204, ssp. Brandegei, 203, ssp. Caseana, 201, 256, ssp. Cusickii, 205, ssp. hastata, 206; cava, 194; chihuahuana, 234; crassipedicellata, 234; crystallina, 217, var. strictissima, 217; Cucullaria, 239; curvisiliqua, 226, ssp. curvisiliqua, 226, ssp. grandibracteata, 227; curvisiliqua var. grandibracteata, 228, var. tenerior, 222; curvisiliquaeformis, 234; Cusickii, 205, var. bastata, 206; densicoma, 230; Engelmannii, 230, var. exaltata, 230; euchlamydea, 230; sect. Eucorydalis, 209; eximia, 239; flavidula, 215; flavula, 215; formosa, 239; fungosa, 239; Geyeri, 215; glauca, 211; Gooddingii, 229; Halei, 222; bastata, 206; Hendersonii, 205; bypecoiformis, 229; idaboensis, 205; isopyroides, 230, var. Mearnsii, 230; Jonesii, 229, var. stenophylla, 234; lutea, 239; macrophylla, 199; macrorrbiza, 229; micrantha, 219, ssp. australis, 222, ssp. micrantha, 219, ssp. texensis, 225; micrantha var. diffusa, 219, var. leptosiliqua, 222, var. pachysiliquosa, 219; monilifera, 229, var. ferruginifera, 219; montana, 234; ochotensis, 239; oregana, 230; pachyloba, 234; paeoniaefolia, 239; pauciflora, 207, 258, var. Chamissonis, 207, var. parviflora, 207; sect. Pes-gallinaceus, 207; pseudomicrantha, 237, var. Griffithsii, 234; pumila, 194; sect. Ramoso-sibiricae, 198; roses, 211; Scouleri, 199, 239, 256; sempervirens, 211; tenuifolia, 239; tortisiliqua, 229, var. longibracteata, 230; washingtoniana, 229; Wetherillii, 229; wyomingensis, 229, var. lativaginata, 230 Corydallis, 197 Cryptoceras, 197 Cucumber, "Small Sugar," 102, 108

Cucumber, "Small Sugar," 102, 108 Cucumer: Meer, 102, 108; Türckisch, 102, 108

Cucurbita, 101; capitata, 102; digitata, 103; ficifolia, 101, 104, 106; indica, 102, III; Lagenaria, 106; maxima, 101, 103, 107, III; melanosperma, 103, 104; mixta, 106; moschata, 101, 102, 104, 105, 106; Pepo, 101, 108, III, varieties of, 102, 106, III; perennis, 103; verrucosa, 102, 106, III

Cucurbits: cultivated, American origin of the, 101; evidence from the herbals, 101; evidence for the New World origin, 105; recent botanical evidence, 106; survey of old and recent botanical evidence, 103

Cuidad Vieja, Guatemala, corn from, 457 Cupressinoxylon sp., 153, transverse and tangential section of, associated with Tempskya, 182

Cushaw, 105, 106

Cycad species, compared with Tempskya Wessellii, 133 Cycadeoidea, 151, 153

Cysticapnos, 188

D

Dianthus caryophyllus, Inheritance of flower color in, 39

Dicentra, 191; canadensis, 239; Cucullaria, 239; eximia, 239; formosa, 239 Dicksonia fibrosa, 131, 142, 145

Dicotyledoneae, fossil, some notes and methods on, 153

Dodge, Carroll W. Mycocandida riboflavina, 31

Dueñas, Guatemala, corn from, 455, 456, 458

Dynamics of leaf variation in Asclepias tuberosa, Some, 353

E

Economic aspects of Hevea, in Peru, 261 Economic importance of species of Corydalis, 195 Ethnobotany of Hevea, 303, of maize, 434

1

Fermentation of galactose and maltose in Saccharomyces, 95 Ferns, fossil, 132 Flower color in carnation, inheritance of, 39; color groups, 41

Fomes, 113; applanatus, 114; idahoensis, fossil, 113, 114; pinicola, 114

Fossil hunter, John Henry Britts, physician and, 115

Fossil plants: The Idaho Tempskyas and associated, 119; collected by Dr. John Henry Britts, 115; collected by C. Henry Thomas, 185

Fossil polypores from Idaho, 113

Fumaria, 188; aurea, 229; flavula, 215; glauca, 211; pauciflora, 207; sempervirens, 211; vesicaria, 188

0

Galactose, ability of yeast cultures to ferment, 96

Geissman, T. A., Gustav A. L. Mehlquist and. Inheritance in the carnation (Dianthus caryophyllus). III. Inheritance of flower color, 39

Genes, organized, versus molecular, 92 Geology of South America, 299

Gleichenites coloradensis, 152

Gourds, 106
Guatemalan maize: Field studies of, 433, 430, 440; Appendix, 453-467; brewing varieties, 445; common varieties, 452-467; compared with Mexican maize, 438; measurements of collections, 437; popping varieties, 445; salpor, 448

Н

Hemitelia crenulata, 141, 144, 178; Smithii, 143, 180

Herbals, evidence from, of American origin of Cucurbita, 101

Herbarium: use of, for plant population studies, 362; profile compared with roadside profile, 403, 407

Hevea, A study of (with its economic aspects) in the Republic of Peru, 261; cytology of, 287; distribution of, in Peru, 309, map showing, 310; ethnobotany of, 303; hybridization of, 286, 305; morphology of, 264, buds and flowers, 280, 330, 334-336, leaflets, 277, short-shoots, 270, 328-332; taxonomy, 292; variation, 287.

Hevea, 292; Benthamiana, 276, 304, 336, 344-348; brasiliensis, 276, 305, 328, 332, 340, 344-348, var. acreana, 306, var. angustifolia, 305, var. cuneata, 294, mut.

Granthami, 266, X guianensis var. lutea, 314, var. janeirensis, 306, var. latifolia, 305, X pauciflora, 317, 352, f. Randiana, 306, var. Randiana, 306, var. stylosa, 305, f. subconcolor, 306, f. typica, 306; camporum, 265; caucho, 292; collina, 292; confusa, 300; cuneata, 294; Foxii, 319; glabrescens, 319, 320; guianensis, 276, 292, 330, 334-348, var. collina, 292, var. cuneata, 292, 294, var. lutea, 293, 334-348, var. marginata, 319, ssp. occidentalis, 292; var. occidentalis, 292, X pauciflora, 318, ssp. typica, 292; guianensis var. lutea, 293, X Benthamiana, 319, X pauciflora, 320, 352, X pauciflora X brasiliensis, 321, f. peruviana, 294; bumilior, 321; janeirensis, 305; Kunthiana, 305; lutea, 293, var. cuneata, 294, f. pilosula, 294, var. pilosula, 294, var. typica, 294; membranacea, 300, f. leiogyne, 300, var. leiogyne, 300; microphylla, 276, 285, 292, 342-350; minor, 265; nigra, 292; nitida, 276, 297, 338, 344-350, var. toxicodendroides, 265; paludosa, 321; pauciflora, 276, 300, 328, 338, 344-348, ssp. coriacea, 300, var. coriacea, 300, ssp. typica, 300; peruviana, 293; Randiana, 305; rigidifolia, 276, 291, 332, 340, 344-350; Sieberi, 305; Spruceana, 276, 285, 291, 330, 336, 344-348; viridis, 297, var. toxicodendroides, 265

Hybrid yeasts, 75

Hybridization: in Asclepias tuberosa, 379; in Hevea, 286

Hybrids, putative, of Hevea in Peru, 313

T

Idaho: Fossil polypores from, 113; northeast corner of Lanes Creek quadrangle, 122, 123; Tempskyas and associated fossil plants from, 119; Some comments on the discovery of Tempskya in the vicinity of Wayan, by C. Henry Thomas, 185; view of hills southeast of Wayan, 158

Inheritance: in the carnation (Dianthus caryophyllus). III. Inheritance of flower color, 39; Mendelian, of genes affecting vitamin-synthesizing ability in Saccharomyces, 95

Insect visitants of Asclepias tuberosa, 356,

J

Jatropha elastica, 292

K

Kern, Ellen M., Henry N. Andrews and. The Idaho Tempskyas and associated fossil plants, 119 Kloeckera (Pseudosaccharomyces), 33

L

Lagenaria, 105; siceraria, 101 Leaf variation in Asclepias tuberosa, 353 Lenz, Lee W., Henry N. Andrews and. Fossil polypores from Idaho, 113

Lindegren, Carl C.: and Gertrude Lindegren. Mendelian inheritance of genes affecting vitamin-synthesizing ability in Saccharomyces, 95; and Caroline Raut. A direct relationship between pantothenate concentration and the time required to induce the production of pantothenate-synthesizing "mutants" in yeasts, 85; The effect of the medium on apparent vitamin-synthesizing ability in Saccharomyces, 75

Lindegren, Gertrude, Carl C. Lindegren and. Mendelian inheritance of genes affecting vitamin-synthesizing deficiencies of microorganisms, 95

Lorain, John, descriptions of early corn varieties by, 15

M

Macock, 105

Maíz chapolote, 44; maíz huajaqueña, 458; maíz negrito, 459

Maize, Guatemalan: Field studies of, 433; see Corn and Zea Mays

Maltose, ability of yeast cultures to ferment, 96

Mass collection studies of Asclepias tuberosa, 363

Media: Burkholder's, 75, growth of yeast cultures on, 76, 81, 85, 95; formulas of different synthetic, 78; Hutner's, 75, growth of yeast cultures on, 76, 82

Medio Monte, Guatemala, corn from, 456 Medium, The effect of, on apparent vitamin-synthesizing deficiencies of microorganisms, 75

Mehlquist, Gustav A. L., and T. A. Geissman. Inheritance in the carnation (Dianthus caryophyllus). III. Inheritance of flower color, 39

Melibiose-fermenting capacity of various yeast cultures, 77

Melons, 104

Melo-pepo, 102; clypeatus, 102; compressus, 102; latior Clypeiformis, 102, III; teres, 102

Mendelian inheritance of genes affecting vitamin-synthesizing ability in Saccharomyces, 95

Microorganisms, The effect of the medium on apparent vitamin-synthesizing deficiencies of, 75

Milkweed, see Asclepias

Missouri, paleobotanical work in, 115 Mixco, Guatemala, corn from, 464

Mutants in yeasts, pantothenate-synthesizing, A direct relationship between pantothenate concentration and the time required to induce the production of, 85

Mylocyprinus robustus, 113 Mycocandida riboflavina, 31, 34, 36

N

Neckeria, 188, 197; aurea, 229, var. occidentalis, 234; curvisiliqua, 226; flavula, 215; glauca, 211; micrantha, 219; sempervirens, 211

Northern flint corns, 1: chromosome knob numbers for, 6, 10, 17; morphology of, 2; origin of, 18; varieties from the northeast, 6, from the Great Plains and Midwest, 7; varieties of: Dakota White, 22, pl. 3, Longfellow Flint, 3, 12, Parker's Flint, 22, pl. 2, Spanish Popcorn, 22, pl. 5, Stevens' Flint, 11, 22, pl. 1, Twelverow Dakota, 22, pl. 4.

0

Odoptera, 197; aurea, 229

Ownbey, Gerald B. Monograph of the North American species of Corydalis, 187

P

Paleobotanical collecting in Missouri, 115, in Idaho, 113, 119

Paleogeography of eastern North America,

Pantothenate: concentration, A direct relationship between, and the time required to induce the production of pantothenate-synthesizing mutants in yeasts, 85; genes affecting synthesis of, 95

Paraminobenzoic acid, genes affecting synthesis of, 97 Paramos, Guatemala, corn from, 463 Patzún, Guatemala, corn from, 439, 452 Pepo, 103; Indicus minor angulosus, 103, III; maximus indicus compressus, 102, III; maximus oblongus, 102; minor clypeatus, 103, minor oblongus, 103; minor rotundus, 103

Pepones lati, 102, III

Peru: A study of Heavea in the Republic of, 261; map of, 310

Phenocontour maps, use of, in study of Asclepias tuberosa, 370 Phoenix dactilifera, 132

Pioneer Hi-Bred Corn Co., experimental plots of, 1

Pistolochia, 197

Polypores, fossil, from Idaho, 113

Popcorn in Guatemala, 454, balls of, 447, being sold in market-place, 439

Population patterns of three subspecies of Asclepias tuberosa, 362

Prehistoric corn collections, map of eastern United States showing, 13

Pseudofumaria, 188 Pseudo-Fumaria, 197 Pumpkin, field, 102

Pyridoxine, genes affecting synthesis of, 97

Q

Quezeltenango, Guatemala, corn from, 452, 453, 466, 467

R

Raut, Caroline, Carl C. Lindegren and. A direct relationship between pantothenate concentration and the time required to induce the production of pantothenate-synthesizing mutants in yeasts, 85; The effect of the medium on the apparent vitamin-synthesizing deficiencies of microorganisms, 75

Roadside profile of butterflyweed, analysis of, 394, compared with herbarium profile,

Rubber, Brazilian, 266; economic importance of, 261; history of, in South America, 261

Saccharomyces: Mendelian inheritance of genes affecting vitamin-synthesizing ability in, 95; carlsbergensis, effect of medium on vitamin synthesis with, 75, 91, pedigree of a hybrid between, and S. cerevisiae, 79; cerevisiae, effect of medium on vitamin synthesis with, 75, 85, hybrid of, heterozygous for mating type and fermentative ability, 96; fragilis, 31, 34

Salcajá, Guatemala, corn from, 454

Salpor in Guatemala, 448 San Antonio Aguascalientes, Guatemala, corn from, 439, 455

San Juan de l'Obispo, Guatemala, corn from, 460

San Lucas, Guatemala, corn from, 461 San Miguel Morazón, Guatemala, corn from, 462

Santa Lucía, Guatemala, corn from, 459 Seibert, R. J. A study of Hevea (with its economic aspects) in the Republic of Peru, 261

Siphonia apiculata, 293; brasiliensis, 305; brevifolia, 293; Cabuchu, 292; elastica, 292; guianensis, 292; Kunthiana, 305; lutea, 293; nitida, 297; pauciflora, 300; ridleyana, 306

Sophorocapnos, 197 Squash, 101: "Acorn," 103; "Golden Custard," 102, 111; "Perfect Gem," 103; "Table Queen," 103, III; Warted, 106; "White Bush Scallop," 102, III Syringospora, 34

T

Tassel branches in northern flint corns, 2, 4 Tempskya, 119; grandis, 146, 147; Knowltoni, 121, 147; minor, 130, 147; rossica, 146, 147; Schimperi, 149; Wessellii, 124, compared with living plants, 133; Whitei, 124; wyomingensis, 124, 147, 151

Tempskya: annuli spores and fragments, 144; associated plant remains, 151; as animal food, 152; compared with other plants, 143; disc specimens, 127, 166, 174; longitudinal sections through, 138; restoration showing probable habit, 140; roots, 139, 168, 170; spores and sporangia, 149; stems, 136, 172-176; taxonomic considerations, 146; transverse sections of specimens, 126, 135, 137, 176; trunks, 125, 131, 160-164

Tempskyas, The Idaho, and associated fossil plants, 119

Thiamin, genes affecting synthesis of, 95 Thomas, Mr. C. Henry: fossil-collecting activities of, 120; with a portion of his Tempskya collection, 158; Some comments on the discovery of Tempskya in the vicinity of Wayan, Idaho, by, 185

Thyrsopteris elegans, 149

Todea barbara, 145 Tripsacum, 443; australe, 19

V Variation: leaf, in Asclépias tuberosa, 353;

pattern of, in Zea Mays, 433
Vegetable marrow, 102
Viability of cells, relation between, and inability to grow on deficient medium, 83
Vitamin-synthesizing: ability in Saccharomyces, Mendelian inheritance of, 95; de-

of the medium on apparent, 75 Vitamins, effect of multiple deficiencies of, on growth of Saccharomyces cerevisiae, 82

ficiencies of microorganisms, The effect

W

Whitaker, Thomas W.: American origin of the cultivated cucurbits, 101 White, Dr. David, 115

Woodson, Robert E., Jr.: Some dynamics of leaf variation in Asclepias tuberosa, 353

Wyoming, fossil plants from, 120

Y

Yeast, A new, 31

Yeasts: A direct relationship between pantothenate concentration and the time required to induce the production of pantothenate-synthesizing mutants in, 85

Yeasts, see Saccharomyces

Z

Zea Mays, see Corn
Zea Mays, variation pattern of, 433
Zuñil, Guatemala, corn from, 440, 452, 464,
465

